

Unification of standard and exotic matter through a $\mathbf{Z}_2$ symmetry

Ernesto A. Matute*

Departamento de Física, Universidad de Santiago de Chile, Casilla 307, Santiago 2, Chile

We consider a scenario in which the discrete weak symmetry between quarks and leptons is extended to the weak force by introducing exotic partners. We conjecture that there exists a hidden discrete symmetry $\tilde{\mathbf{P}}$, defining a $\mathbf{Z}_2$ group, between standard and exotic quarks and leptons. The unified model $\text{SU}(3)_q \times \text{SU}(3)_{\tilde{q}} \times \text{SU}(2)_{q\tilde{\ell}} \times \text{SU}(2)_{\tilde{q}\ell} \times \text{U}(1)_Y \times \tilde{\mathbf{P}}$ is discussed, where the unifying discrete symmetry extends over particles and forces. It is shown that the lighter neutral and charged weak bosons generated upon spontaneous symmetry breaking have the same properties as those of the standard model. Cabbibo-Kobayashi-Maskawa unitarity is used to set a mass of order 2.8 TeV for the nonstandard weak bosons, which do not exhibit quark-lepton universality. A grand unified theory of $(\text{GUT})_{q\tilde{\ell}} \times (\text{GUT})_{\tilde{q}\ell} \times \tilde{\mathbf{P}}$ type, with decay of exotic matter into standard matter and no decay of the ordinary nucleon, is put forward.

PACS numbers: 12.60.Cn; 11.30.Hv; 12.10.Dm; 14.70.Pw

The unification of quark and lepton properties under strong and electroweak interactions through a continuous or discrete symmetry has been one of the targets of research efforts over the last few decades [1]. Mainly the idea has been to assign similar degrees of freedom to quarks and leptons, extending in a sense the color gauge group of quarks to include leptons. Nonstandard or exotic gauge bosons and fermions are commonly added to the spectrum of the standard model (SM). The purpose of this paper is to implement these developments with the notion that there is a hidden $\mathbf{Z}_2$ symmetry in nature unifying the properties of standard and exotic fermions. And that these interact with separate but symmetric gauge bosons, so duplicating the gauge group $\text{SU}(3)_q \times \text{SU}(2)_L \times \text{U}(1)_Y$ of the SM and its unified extensions. In the scenario of a duplicated grand unified theory (GUT) with such a discrete symmetry, the decay of exotic matter into standard matter and the stability of the ordinary nucleon are expected. This makes an interesting point as nucleon decay is one of the pillars of GUTs and no ordinary proton decay has yet been observed.

In the strong interaction sector of the SM there are substantial differences between quarks and leptons as they come in triplets and singlets of the color gauge group $\text{SU}(3)_q$, respectively. However, in the electroweak sector their properties are so similar that it is conceivable an interaction gauge symmetry completely or at least partially duplicated in the form $G_q \times G_\ell$ but unified through a discrete symmetry; the subscripts q and ℓ are to emphasize that each G only acts on quarks or leptons. A trouble with this picture is that the $\text{U}(1)_Y$ charges of quarks and leptons are different. Nonetheless, the symmetric duplication of the electroweak gauge group produces triangle anomalies which can be cancelled by introducing exotic quarks and leptons, related to the standard ones through the same discrete symmetry. This pattern then allows to expand the standard electroweak gauge group to the

unified form $G_{q\tilde{\ell}} \times G_{\tilde{q}\ell} \times \tilde{\mathbf{P}}$, where $\tilde{q}$ and $\tilde{\ell}$ denote exotic quarks and exotic leptons, respectively, and $\tilde{\mathbf{P}}$, hereafter referred to as exotic symmetry, denotes the discrete symmetry which transforms standard quarks and leptons to exotic ones and vice versa: $q \leftrightarrow \tilde{q}$, $\ell \leftrightarrow \tilde{\ell}$. This unifying symmetry defines a $\mathbf{Z}_2$ group as $\tilde{\mathbf{P}}^2 = 1$. Moreover, this discrete exotic symmetry is extendible to strong interactions.

Our aim in this paper is to discuss the unified gauge group $\text{SU}(3)_q \times \text{SU}(3)_{\tilde{q}} \times \text{SU}(2)_{q\tilde{\ell}} \times \text{SU}(2)_{\tilde{q}\ell} \times \text{U}(1)_Y \times \tilde{\mathbf{P}}$, assuming that each set of fermions is separately conserved. It is a unified model in the sense that the duplicated gauge group factors have the same gauge coupling constant, and the standard and exotic quarks and leptons have similar quantum numbers. Symmetric standard and exotic quarks interact with symmetric gluons and weak bosons. The two $\text{SU}(3)$ color groups are unbroken and exotic gluons confine exotic quarks into heavy exotic hadrons. Any explanation for the mass hierarchy of standard quarks should be extended to include exotic quarks, since standard and exotic gluons confine at the same scale; in a sense, three new heavy generations of quarks and leptons are added to the standard spectrum. We may say that the exotic symmetry $\tilde{\mathbf{P}}$ has the incentive of expanding the electroweak quark-lepton symmetry to the strong interaction sector. We note here that the spontaneous breakdown of the discrete symmetry $\tilde{\mathbf{P}}$ leads to well-known cosmological problems associated with topological defects, namely, domain walls; they can be solved, however, via interactions of domain walls with primordial black holes [2]. The model provides a framework for understanding possible deviations from the SM predictions. We use the reported quark-lepton universality violation of weak interactions at low energies to fix the scale of the new weak physics.

Specifically, the transformation properties of standard quarks and leptons are

$$\begin{aligned} q_{nL}^i &\sim (3, 1, 2, 1, \frac{1}{3}), & \ell_{nL} &\sim (1, 1, 1, 2, -1), \\ u_{nR}^i &\sim (3, 1, 1, 1, \frac{4}{3}), & \nu_{nR} &\sim (1, 1, 1, 1, 0), \end{aligned}$$

*Electronic address: ematute@lauca.usach.cl (E.A. Matute).

$$d_{nR}^i \sim (3, 1, 1, 1, -\frac{2}{3}), \quad e_{nR} \sim (1, 1, 1, 1, -2), \quad (1)$$

where n denotes the three generations, $q_{nL} = (u_{nL}, d_{nL})$ and $\ell_{nL} = (\nu_{nL}, e_{nL})$ are the $SU(2)$ doublets, i denotes the color degree of freedom, and the numbers in brackets describe the transformation properties under the unified gauge group. The ν_R 's are introduced because of the experimental signatures for neutrino masses [3].

The quark and lepton assignments in Eq. (1), however, introduce axial anomalies of the $U(1)_Y[SU(2)]^2$ type. They are associated with the nonvanishing trace of Y for quarks and for leptons due to the separation of the weak gauge group. To achieve their cancellation we include exotic fermions as done in the so-called ununified models [4]. We consider the exotic quarks and leptons

$$\begin{aligned} \tilde{q}_{nL}^i &\sim (1, 3, 1, 2, \frac{1}{3}), & \tilde{\ell}_{nL} &\sim (1, 1, 2, 1, -1), \\ \tilde{u}_{nR}^i &\sim (1, 3, 1, 1, \frac{4}{3}), & \tilde{\nu}_{nR} &\sim (1, 1, 1, 1, 0), \\ \tilde{d}_{nR}^i &\sim (1, 3, 1, 1, -\frac{2}{3}), & \tilde{e}_{nR} &\sim (1, 1, 1, 1, -2). \end{aligned} \quad (2)$$

Under the exotic symmetry $\tilde{P}$, ordinary and exotic fermions are transformed as follows:

$$\begin{aligned} q_{nL}^i &\leftrightarrow \tilde{q}_{nL}^i, & u_{nR}^i &\leftrightarrow \tilde{u}_{nR}^i, & d_{nR}^i &\leftrightarrow \tilde{d}_{nR}^i, \\ \ell_{nL} &\leftrightarrow \tilde{\ell}_{nL}, & \tilde{\nu}_{nR} &\leftrightarrow \nu_{nR}, & \tilde{e}_{nR} &\leftrightarrow e_{nR}, \end{aligned} \quad (3)$$

and $W_q^a \leftrightarrow W_{\tilde{q}}^a$, $G_q^b \leftrightarrow G_{\tilde{q}}^b$, where W_q^a , $W_{\tilde{q}}^a$ are the weak gauge bosons of $SU(2)_{q\tilde{\ell}}$, $SU(2)_{\tilde{q}\ell}$, and G_q^b , $G_{\tilde{q}}^b$ are the gluons of $SU(3)_q$, $SU(3)_{\tilde{q}}$, respectively. Besides, the corresponding bare gauge couplings are required to be equal. Thus the whole Lagrangian of electroweak and strong interactions, with the Higgs fields defined below, becomes invariant under $\tilde{P}$ symmetry. This leads to a great simplification of many results which cannot be achieved for ununified models due to current structures; phenomenological implications are therefore different.

In this work we assume that the weak gauge symmetry and exotic symmetry $\tilde{P}$ are spontaneously broken at one stage, which happens with our choice of Higgs multiplets. The minimal set of colorless scalar multiplets that breaks the electroweak gauge symmetry $SU(2)_{q\tilde{\ell}} \times SU(2)_{\tilde{q}\ell} \times U(1)_Y \times \tilde{P}$ down to $U(1)_{em}$ and gives mass to all fermions of the model includes

$$\begin{aligned} \phi_q &\sim (2, 1, +1), & \phi_{\tilde{q}} &\sim (1, 2, +1), \\ \Sigma &= \sigma + i\boldsymbol{\tau} \cdot \boldsymbol{\pi} \sim (2, 2, 0), \end{aligned} \quad (4)$$

with vacuum expectation values (VEVs)

$$\begin{aligned} \langle \phi_q \rangle &= \begin{pmatrix} 0 \\ v/\sqrt{2} \end{pmatrix}, & \langle \phi_{\tilde{q}} \rangle &= \begin{pmatrix} 0 \\ v/\sqrt{2} \end{pmatrix}, \\ \langle \Sigma \rangle &= \begin{pmatrix} u & 0 \\ 0 & u \end{pmatrix}. \end{aligned} \quad (5)$$

The Pauli matrices $\boldsymbol{\tau}$ in Eq. (4) belong to the standard gauge group $SU(2)_L$ and they are there to indicate that

after spontaneous symmetry breaking Σ splits into a singlet and a triplet. Each row and each column of Σ is a doublet under $SU(2)_{q\tilde{\ell}}$ and $SU(2)_{\tilde{q}\ell}$, respectively. Under $\tilde{P}$ symmetry $\phi_q \leftrightarrow \phi_{\tilde{q}}$ and $\Sigma \leftrightarrow \Sigma$. The equality of the bare VEVs of ϕ_q and $\phi_{\tilde{q}}$ is broken by radiative corrections due to the hierarchy of Yukawa couplings responsible for fermion masses, so that the renormalized quantities are actually equal up to finite radiative corrections. We remark that this type of Higgs multiplets is also present in ununified models, but without any restriction on the VEVs of Higgs doublets.

The gauge symmetry breaking follows the pattern

$$\begin{aligned} &SU(2)_{q\tilde{\ell}} \times SU(2)_{\tilde{q}\ell} \times U(1)_Y \times \tilde{P} \\ &\quad \downarrow \\ &SU(2)_L \times U(1)_Y \\ &\quad \downarrow \\ &U(1)_{em}. \end{aligned} \quad (6)$$

The unbroken electric charge generator is given by

$$Q = I_{3L} + \frac{1}{2}Y = I_{3q} + I_{3\tilde{q}} + \frac{1}{2}Y, \quad (7)$$

and the electroweak couplings of the model, in terms of the standard electromagnetic coupling constant and Weinberg angle, take the form

$$g = g_q = g_{\tilde{q}} = \frac{\sqrt{2}e}{\sin \theta_W}, \quad g' = \frac{e}{\cos \theta_W}. \quad (8)$$

The two physical sets of left-handed weak bosons and the photon get the following linear combinations of the electroweak gauge bosons W_q , $W_{\tilde{q}}$ and A :

$$\begin{aligned} W_1^\pm &= \frac{1}{\sqrt{2}}(W_q^\pm + W_{\tilde{q}}^\pm), \\ Z_1 &= \frac{1}{\sqrt{2}}\cos \theta_W(W_q^3 + W_{\tilde{q}}^3) - \sin \theta_W A, \\ A_{em} &= \frac{1}{\sqrt{2}}\sin \theta_W(W_q^3 + W_{\tilde{q}}^3) + \cos \theta_W A, \\ W_2^\pm &= \frac{1}{\sqrt{2}}(W_q^\pm - W_{\tilde{q}}^\pm), \\ Z_2 &= \frac{1}{\sqrt{2}}(W_q^3 - W_{\tilde{q}}^3). \end{aligned} \quad (9)$$

They pick up the mass

$$\begin{aligned} m_{W_1}^2 &= \frac{1}{4}g^2v^2, & m_{Z_1}^2 &= \frac{1}{4}(g^2 + 2g'^2)v^2 = \frac{m_{W_1}^2}{\cos^2 \theta_W}, \\ m_{A_{em}} &= 0, & m_{W_2}^2 &= \frac{1}{4}g^2(v^2 + 2u^2), \\ m_{Z_2} &= m_{W_2}. \end{aligned} \quad (10)$$

The gauge fields $\mathbf{W}_1 = (\mathbf{W}_q + \mathbf{W}_{\tilde{q}})/\sqrt{2}$ are identified with the SM weak bosons, while the orthogonal combination $\mathbf{W}_2 = (\mathbf{W}_q - \mathbf{W}_{\tilde{q}})/\sqrt{2}$ represents the nonstandard ones. We note that the standard mass relation

$m_{W_1} = m_{Z_1} \cos \theta_W$ is preserved. Also significant, the heavier nonstandard neutral and charged weak bosons acquire the same mass. All of these only occurs within the context of the unified model of ordinary and exotic quarks and leptons in which the exotic symmetry $\tilde{P}$ is essential, making the difference from other extensions of the SM.

The charged-current interactions involving the $W_1^\pm$ and $W_2^\pm$ left-handed bosons can be written as

$$\mathcal{L}_{CC} = \frac{g}{2}(W_{1\mu}^+ J_{1L}^{-\mu} + W_{2\mu}^+ J_{2L}^{-\mu} + \text{h.c.}), \quad (11)$$

with weak currents defined, at the ordinary and exotic quark and lepton level, by

$$\begin{aligned} J_{1L}^\pm &= J_{qL}^\pm + J_{\ell L}^\pm + J_{\tilde{\ell} L}^\pm + J_{\tilde{q} L}^\pm, \\ J_{2L}^\pm &= J_{qL}^\pm - J_{\ell L}^\pm + J_{\tilde{\ell} L}^\pm - J_{\tilde{q} L}^\pm. \end{aligned} \quad (12)$$

As for the neutral-current interactions involving the Z_1 and Z_2 left-handed bosons, we have

$$\begin{aligned} \mathcal{L}_{NC} &= \frac{g}{\sqrt{2} \cos \theta_W} Z_{1\mu} (J_{1L}^{o\mu} - \sin^2 \theta_W J_{em}^\mu) \\ &\quad + \frac{g}{\sqrt{2}} Z_{2\mu} J_{2L}^{o\mu}, \end{aligned} \quad (13)$$

where now

$$\begin{aligned} J_{1L}^o &= J_{qL}^3 + J_{\ell L}^3 + J_{\tilde{\ell} L}^3 + J_{\tilde{q} L}^3, \\ J_{2L}^o &= J_{qL}^3 - J_{\ell L}^3 + J_{\tilde{\ell} L}^3 - J_{\tilde{q} L}^3, \\ J_{em} &= J_{qL}^3 + J_{\ell L}^3 + J_{\tilde{\ell} L}^3 + J_{\tilde{q} L}^3 \\ &\quad + J_{qY} + J_{\ell Y} + J_{\tilde{\ell} Y} + J_{\tilde{q} Y}. \end{aligned} \quad (14)$$

We see, from Eqs. (8)–(14), that $W_1^\pm$ and Z_1 have the SM mass and coupling relationships with a universality that comprises exotic fermions, and no mixing with the nonstandard $W_2^\pm$ and Z_2 exists, which exhibit no quark-lepton universality and also the breaking of exotic symmetry. At the mass scale of the SM, this corresponds to adding three new generations of quarks and leptons.

We next write the most general renormalizable, gauge invariant and $\tilde{P}$ symmetric Lagrangian for the Yukawa couplings of Higgs bosons with fermions, consistent with separate ordinary and exotic baryon number conservation, as well as separate ordinary and exotic lepton number conservation, which are associated with $U(1)$ global vector symmetries:

$$\begin{aligned} \mathcal{L}_{YUK} &= G_{nm}^d \left(\bar{q}_{nL} \phi_q d_{mR} + \bar{\tilde{q}}_{nL} \phi_{\tilde{q}} \tilde{d}_{mR} \right) \\ &\quad + G_{nm}^u \left(\bar{q}_{nL} \phi_q^c u_{mR} + \bar{\tilde{q}}_{nL} \phi_{\tilde{q}}^c \tilde{u}_{mR} \right) \\ &\quad + G_{nm}^e \left(\bar{\ell}_{nL} \phi_{\tilde{q}} e_{mR} + \bar{\tilde{\ell}}_{nL} \phi_q \tilde{e}_{mR} \right) \\ &\quad + G_{nm}^\nu \left(\bar{\ell}_{nL} \phi_q^c \nu_{mR} + \bar{\tilde{\ell}}_{nL} \phi_{\tilde{q}}^c \tilde{\nu}_{mR} \right) \\ &\quad + \text{h.c.} \end{aligned} \quad (15)$$

Because of gauge invariance, only the ordinary or exotic lepton number conservation is related to an apparently *ad*

hoc global symmetry. The left-handed bidoublet Σ then has no Yukawa couplings to fermions. Substitution of VEVs leads to mass matrices that can be made real and diagonal by suitable unitary transformations of the left- and right-handed components of the pertinent fermion fields. In this mass eigenstate basis, Z_1 and Z_2 have flavor-conserving couplings.

The equality between Yukawa couplings of ordinary and exotic particles yields, however, equal mass values at tree level. These mass relationships can easily be avoided if two new Higgs doublets $\phi'_q \sim (2, 1, -1)$ and $\phi'_{\tilde{q}} \sim (1, 2, -1)$ are added, which under $\tilde{P}$ symmetry transform as $\phi'_q \leftrightarrow \phi_q^c$ and $\phi'_{\tilde{q}} \leftrightarrow \phi_{\tilde{q}}^c$. It can be seen that in this case the new Yukawa Lagrangian leads to no standard and exotic mass relations if the VEVs of the new Higgs doublets are equal to each other but different from those in Eq. (5). This complication in the Higgs sector has no effect on the properties of the gauge bosons discussed in this paper, except that their masses now receive contributions from the new doublets (see Eq. (10)). As mentioned above, all of this happens because of the $\tilde{P}$ symmetry for the full Lagrangian. Clearly, this scenario, rendering the model rather uneconomic but richer phenomenologically, favors the SM with two Higgs doublets.

Unfortunately, the systematics of the masses of the known quarks and leptons are unexplained, so that there is no reliable extrapolation to masses of exotic generations. However, strict constraints on the masses of the new fermions are obtained from bounds placed by cosmology and precision electroweak experiments.

On the one side, cosmological evidences on the ordinary baryon asymmetry and absence at present of exotic baryons made of exotic quarks require that exotic matter be unstable, decaying rapidly enough into observable ordinary matter [3]. As argued below, this is a strong motivation to embed the duplicated SM into a duplicated GUT such that exotic quarks and ordinary leptons transform under one GUT gauge group, and exotic leptons and ordinary quarks do under the other. Besides of allowing the decay of exotic matter, such a duplicated GUT would indicate the ordinary proton stability. In this scenario, the cosmological ordinary baryon (lepton) asymmetry requires the exotic lepton (baryon) asymmetry. Thus, the problem of ordinary baryogenesis (leptogenesis) is also that of exotic leptogenesis (baryogenesis) [5].

On the other side, there are constraints on the masses and mass-splittings of exotic fermions from the standard Z and W gauge boson observables. If any of the exotic fermions is lighter than about half of the mass of the SM-like Z boson of the model, this will affect its decay width very strongly. For instance, if the Z can decay into exotic light neutrinos, its width will be significantly increased, in contradiction with experiment. In fact, the direct measurement of the invisible Z width gives the bound $N_\nu = 2.92 \pm 0.07$ on the number of light neutrino types [3]. On the contrary, if the exotic fermions are heavy, their couplings to the lighter Z and W bosons will affect the values of the precision electroweak

observables S , T and U , or equivalent oblique parameters on the gauge self-energies at the loop level [6]. These parameters are defined to constrain heavy new physics, vanishing for the SM. In the case of new generations of fermions, these values restrain the masses and mass-splittings of such fermions. For completeness, we present in the following the contributions to these oblique parameters from each nondegenerate doublet (f_1, f_2) of extra SM-like heavy quarks or leptons with masses (m_1, m_2) , in the limit $m_Z, \Delta m = |m_1 - m_2| \ll m_{1,2}$ [6]:

$$\begin{aligned} S &\simeq \frac{N_c}{6\pi}, \\ T &\simeq \frac{N_c}{12\pi \sin^2 \theta_W \cos^2 \theta_W} \frac{\Delta m^2}{m_Z^2}, \\ U &\simeq \frac{2N_c}{15\pi} \frac{\Delta m^2}{m_1^2}, \end{aligned} \quad (16)$$

where $N_c = 3$ (1) denotes the color number of extra quarks (leptons). We note that if m_1 and m_2 are non-degenerate and not too large, additional negative corrections to S may arise; this parameter in a sense measures the size of the new fermion sector, while T and U do on its weak-isospin breaking. The experimental values for these observables are [3]

$$\begin{aligned} S &= -0.13 \pm 0.10 \text{ } (-0.08), \\ T &= -0.17 \pm 0.12 \text{ } (+0.09), \\ U &= 0.22 \pm 0.13 \text{ } (+0.01), \end{aligned} \quad (17)$$

where the central values assume $m_H=117$ GeV for the SM Higgs boson, and the values shown in the parentheses correspond to the changes for $m_H=300$ GeV; the S and T parameters are highly correlated. It has been shown that three generations of relatively heavy extra quarks and leptons cannot fit the data with the SM Higgs doublet, but they do it with a two-Higgs-doublet extension, although the parameter space becomes very restrictive with fermion masses significantly below 1 TeV [7]; the nonstandard gauge bosons would have an effect on the oblique parameters only at the two-loop level through their couplings to fermions, because as exhibited in Eq. (9) they do not mix with the standard ones [8]. This motivates even more, within the context of our model, the duplication of the SM forces with a two-Higgs-doublet extension included.

Hence, all of these can be met if the exotic fermions are massive and nondegenerate enough to decay into ordinary fermions in a duplicated GUT scenario, with a generic lower mass bound about the standard $m_Z/2$. By contrast, if the masses are significantly greater than the generic bound $\langle H \rangle \simeq 174$ GeV, the Higgs VEV causing the SM-like electroweak symmetry breaking, then the weak interaction among heavy fermions becomes strong and perturbation theory breaks down, so eluding any mass bound depending on the perturbative assumption. Collider searches for new long-lived up and down quarks give by now the lower mass bounds of 220 GeV and 190

GeV, respectively, while the current direct lower bounds on extra stable neutral and charged leptons are about 45.0 GeV and 102.6 GeV, respectively [3]. Moreover, the ordinary and exotic fermions do couple to the degenerate heavy nonstandard Z and W gauge bosons, so that the new physics cannot be fully parametrized in the usual S , T and U framework. Actually, we will use the mass ratio between the light and heavy W (see Eq. (19) below). And to restrain this new parameter, we will consider in this paper charged-current and neutral-current data at the standard matter level.

The relevant charged-current effective Lagrangian for the low-energy ordinary quark-lepton physics becomes

$$\mathcal{L}_{\text{CC}}^{\text{eff}} = \frac{4G_F}{\sqrt{2}(1+\eta)} (J_{1L\mu}^+ J_{1L}^{-\mu} + \eta J_{2L\mu}^+ J_{2L}^{-\mu}), \quad (18)$$

where $J_{1,2L}^\pm$ are the currents in Eq. (12) limited to standard quarks and leptons, η is a parameter defined in the minimal model by

$$\eta = \frac{m_{W_1}^2}{m_{W_2}^2} = \left(1 + \frac{2u^2}{v^2}\right)^{-1}, \quad (19)$$

and G_F is the Fermi constant from the muon decay determined (at tree level) by

$$\frac{4G_F}{\sqrt{2}} = \frac{g^2}{4m_{W_1}^2} (1+\eta). \quad (20)$$

According to Eq. (18), there is no universality in the strength of leptonic and hadronic weak interactions, which depends on the magnitude of the η -parameter. The corresponding effective neutral-current weak interactions are given by an expression similar to Eq. (18), which is derived from Eq. (13) restricted to conventional quarks and leptons.

Therefore, within our scenario the electroweak predictions of the SM also get corrections parameterized by the quantity η defined in Eq. (19). To exhibit the degree of violation of the universality of weak interactions, we consider the constraint dictated by low-energy charged-current data. At the low-energy level of ordinary quarks and leptons, charged-current interactions are governed by the effective Lagrangian of Eq. (18), which can be rewritten as follows:

$$\begin{aligned} \mathcal{L}_{\text{CC}}^{\text{eff}} &= \frac{4G_F}{\sqrt{2}} \left[J_{qL\mu}^+ J_{qL}^{-\mu} + J_{\ell L\mu}^+ J_{\ell L}^{-\mu} \right. \\ &\quad \left. + \frac{1-\eta}{1+\eta} (J_{qL\mu}^+ J_{\ell L}^{-\mu} + J_{\ell L\mu}^+ J_{qL}^{-\mu}) \right]. \end{aligned} \quad (21)$$

Thus, η makes the strength of the leptonic charged-current weak interactions different from the strength of the semileptonic weak interactions, so that semileptonic processes can be considered to constrain its value. We will do this by adapting a calculation by Rajpoot [4] based on preserving Cabbibo universality and Cabbibo-Kobayashi-Maskawa (CKM) unitarity. The relevant four

fermion effective Lagrangian is given by

$$\mathcal{L}_{\text{CC}}^{\text{eff}}(\nu q) = \frac{G_F}{\sqrt{2}} \frac{1-\eta}{1+\eta} V_{qq'} \bar{q}' \gamma_\mu (1+\gamma_5) q \bar{\ell} \gamma^\mu (1+\gamma_5) \nu, \quad (22)$$

where $V_{qq'}$ are the elements of the CKM mixing matrix. Typically these are determined from leptonic decays of hadrons. The data from high precision measurements yields [3]

$$\begin{aligned} \frac{1-\eta}{1+\eta} |V_{ud}| &= 0.9738 \pm 0.0005, \\ \frac{1-\eta}{1+\eta} |V_{us}| &= 0.2200 \pm 0.0026, \\ \frac{1-\eta}{1+\eta} |V_{ub}| &= 0.00367 \pm 0.00047. \end{aligned} \quad (23)$$

Unitarity of the mixing matrix requires $|V_{ud}|^2 + |V_{us}|^2 + |V_{ub}|^2 = 1$. This constraint immediately implies $\eta = 0.00083 \pm 0.00037$ and $m_{W_2} = 2.8 \pm 0.9$ TeV, although the experimental results quoted in Eq. (23) are still controversial [9]. It is worth emphasizing that there is no analogous result in ununified models because the low-energy charged-current structure differs from that of the unified variant.

Additional limits can be obtained from constraints at higher energies coming from precision electroweak measurements. With the increasing accuracy of data, a very convenient choice of free parameters to use in the gauge sector of the extended electroweak model is given by α_{em} , G_F , m_{Z_1} and m_{W_1} , since m_{Z_1} and m_{W_1} are directly measurable. Then the other parameters of the theory like $\sin^2 \theta_W$ and η become dependent parameters defined through Eqs. (8), (10), and (20), by

$$\begin{aligned} \sin^2 \theta_W &= 1 - \frac{m_{W_1}^2}{m_{Z_1}^2}, \\ 1 + \eta &= \frac{\sqrt{2} G_F}{\pi \alpha_{em}} m_{W_1}^2 \sin^2 \theta_W (1 - \Delta r), \end{aligned} \quad (24)$$

where the weak radiative correction parameter Δr has been included. Considering that the SM value of Δr does not change at the energy scale of the model which produces $\eta > 0$, η and therefore m_{Z_2} ($= m_{W_2}$) can be predicted from m_{Z_1} and m_{W_1} . However, they are dominated by the experimental bar errors. We only get an upper bound $\eta < 0.0024$ which implies a lower mass bound of $m_{Z_2(W_2)} > 1.6$ TeV for the nonstandard weak bosons, consistent with the above results. The magnitude of $\Delta r = 0.03434 \mp 0.0017 \pm 0.00014$ used to obtain these limits was extracted from the minimal SM in the on-shell scheme [3]. All nonstandard particles were not included in the radiative corrections predicted by the SM because we search for small tree-level effects of the extra gauge bosons.

Another limit can be derived from Eq. (24) by setting in the SM prediction for m_{W_1} instead of the Δr term.

We approximately find

$$\eta = 2 \frac{\Delta m_{W_1}}{m_{W_1}} \left(\frac{1}{\sin^2 \theta_W} - 2 \right), \quad (25)$$

where $\Delta m_{W_1} = m_{W_1}^{SM} - m_{W_1}$, obtaining forthwith the bounds $\eta < 0.0013$ and $m_{Z_2(W_2)} > 2.2$ TeV. Equation (25) shows how, in the case of the W_1 -mass, a small deviation of the experimental result from the SM prediction defines the η -parameter of the new physics implied by the $\tilde{P}$ -symmetric model. The model favors an experimental value of m_{W_1} below the SM prediction. Lower mass bounds coming from other observables can be computed as well.

Thus, more experimental accuracy is needed to have a signal of the new physics from precision electroweak measurements at high energies, where the SM exhibits a very good agreement with data. We know that the SM cannot be complete, so eventually quantitative disagreements between data and the SM predictions shall occur. We expect that all the magnitudes will be accounted for in part by a simple, consistent value of η , which in turn will give the mass scale for the new left-handed weak bosons. In a sense, the η parameter plays a role similar to that of the mixing angle θ_W in the SM prior to the LEP era; its value obtained from a fit to neutral-current data made possible to predict the mass of weak bosons.

In conclusion, we have analyzed the unified model $\text{SU}(3)_q \times \text{SU}(3)_{\bar{q}} \times \text{SU}(2)_{q\bar{\ell}} \times \text{SU}(2)_{\bar{q}\ell} \times \text{U}(1)_Y \times \tilde{P}$ in which the $\mathbf{Z}_2$ exotic symmetry defined by $\tilde{P}$ is spontaneously broken. It may be viewed as an important stage in the process of full understanding which can be tested at relatively low energies. Our aim in this paper has been limited to account for the observed universality violation of weak interactions and consider phenomenological constraints on the masses and mass-splittings of exotic generations of quarks and leptons that seem to enforce the SM with two Higgs doublets as well as further gauge extensions of our model. What is interesting is that the lighter weak bosons that arise have the same properties as the standard ones, though all predictions of the SM (starting at the tree level) are corrected by η , the new weak parameter introduced by the symmetric model. The magnitude of η actually measures the no universality of weak interactions. We have fixed the value of this parameter using universality/unitarity constraints, which sets a mass of order 2.8 TeV for the nonstandard weak bosons, even though the experimental results used are still controversial. Their existence mean to touch the idea of symmetric left-handed weak forces with experiment, stressing the open question which inquires for the new dynamics that may be behind or associated with exotic symmetry. This work shows, however, that despite the lack of knowledge of this dynamics, much information can be obtained from this symmetry.

As a way forward, the notion that exotic symmetry extends to the forces of the SM already suggests to include the $\text{U}(1)_Y$ interactions to have the unified gauge

group $G_{q\bar{\ell}} \times G_{\bar{q}\ell} \times \tilde{P}$ with $G = SU(3) \times SU(2) \times U(1)$. Next, it is natural to embed each G in a GUT to have $(GUT)_{q\bar{\ell}} \times (GUT)_{\bar{q}\ell} \times \tilde{P}$ which unifies standard and exotic quarks and leptons, and also the strong and electroweak interactions; the $\tilde{P}$ symmetry should be preserved down to energies where the spontaneous breaking of the electroweak gauge symmetry takes place. The ordinary nucleon does not decay in models of this type because of the separation of GUT for ordinary quarks and leptons. Thus, applications of GUT techniques [1] predict transitions between exotic and ordinary quarks and leptons, giving rise to the decay of heavy exotic matter into ordinary matter, with separate conservation of ordinary and exotic color charge. From a cosmological point of view, this is a mandatory condition not to conflict with observational evidences on the absence of exotic baryons. On

the other hand, the well-known cosmological domain wall problems associated with the spontaneous breakdown of the discrete symmetry $\tilde{P}$ can be solved with the mechanism of destroying domain walls by primordial black holes [2], without constraints on the parameters in the Lagrangian of the model. Finally, from the viewpoint of precision electroweak experiments, current constraints on the masses and mass-splittings of new fermions also compel, within the context of our model, the duplication of the SM forces with a two-Higgs-doublet extension included.

This work was partially supported by the Departamento de Investigaciones Científicas y Tecnológicas, Universidad de Santiago de Chile.

-
- [1] For a recent overview, see, e.g., R.N. Mohapatra, *Unification and Supersymmetry* (Springer, New York, 2003), 3rd ed..
 - [2] D. Stojkovic, K. Freese, and G.D. Starkman, Phys. Rev. D **72**, 045012 (2005), and references therein.
 - [3] S. Eidelman *et al.* (Particle Data Group), Phys. Lett. B **592**, 1 (2004).
 - [4] H. Georgi, E.E. Jenkins, and E.H. Simmons, Phys. Rev. Lett. **62**, 2789 (1989); Nucl. Phys. **B331**, 541 (1990); V. Barger and T. Rizzo, Phys. Rev. D **41**, 946 (1990); S. Rajpoot, Phys. Rev. D **41**, 995 (1990); **41**, 1706 (1990); D. Choudhury, Mod. Phys. Lett. A **6**, 1185 (1991); X. Li and E. Ma, Mod. Phys. Lett. A **18**, 1367 (2003).
 - [5] For a review, see W. Buchmüller, hep-ph/0107153, and references therein.
 - [6] M.E. Peskin and T. Takeuchi, Phys. Rev. D **46**, 381 (1992), and references therein.
 - [7] H.J. He, N. Polonsky, and S. Su, Phys. Rev. D **64**, 053004 (2001).
 - [8] M.E. Peskin and J.D. Wells, Phys. Rev. D **64**, 093003 (2001).
 - [9] A. Czarnecki, W.J. Marciano, and A. Sirlin, Phys. Rev. D **70**, 093006 (2004), and references therein.